


View this article online at: patient.info/chalazion

Chalazion

A **chalazion** is a common condition affecting the eyelid and is caused by a blockage of a gland in the eyelid. The swelling (cyst) is usually felt as a small lump. Rarely, it can become infected. If it is causing problems and does not settle on its own, it can be treated with an injection or removed with a small operation.

What is a chalazion?



A chalazion is a small (2-8 mm) fluid-filled swelling (cyst) in the eyelid. It is common and sometimes called a meibomian cyst or tarsal cyst. A chalazion is more common on the upper eyelid. It is possible to have several at once, in more than one eyelid. It is not the same as a stye.

What causes a chalazion?

There are tiny glands just under the inner surface of the eyelid, called meibomian glands. These make an oily fluid called meibum to help lubricate the eye. If the gland becomes blocked then the meibum cannot escape into the tears. It may expand into a swelling (cyst) and leak into the eyelid tissue. This becomes inflamed. Over time, inflammation causes a lump (or granuloma) to form in the eyelid. This is a firm, solid lump that remains for a long time. This lump is the chalazion.

Chalazion is more common in people who have the eye condition blepharitis, or skin conditions such as eczema, because in these conditions the meibum tends to be thicker so the gland is more easily blocked. See separate leaflets called [Stye](#) and [Blepharitis](#) for more details.

What are the symptoms of a chalazion?

- The usual symptom is a small lump which develops on an eyelid.
- Sometimes it causes mild pain or irritation, particularly if it has just started - this usually settles.
- Rarely, it gets infected. It then becomes more swollen, red and painful.
- Sight is not affected, although very occasionally it can become big enough to press on the eyeball and distort vision.

What is the treatment for a chalazion?

- **No treatment** may be necessary as up to half of people get better without any treatment. This can take between two and six months but if the chalazion is not causing you any problems, it is probably best just to watch and wait.
- **Hot compresses** help to ease discomfort. Hold a clean flannel that has been in hot water gently but firmly against the closed eye for 5-10 minutes, 3-4 times a day. Sometimes this warmth and slight pressure is enough to soften the contents of the fluid-filled swelling (cyst), helping it drain more easily. (The water should be hot but comfortable and not scalding.)

- **Antibiotic ointments, drops and medicines** are not recommended as they do not make any difference - the contents of the cyst are infection-free (sterile).
- **Massage** of the cyst after using a hot compress can encourage the cyst to drain. Do this gently, with a clean finger or cotton bud, in the direction of the eyelashes.
- **Cleaning the eyelid** twice per day removes grease and dead skin cells that may contribute to cysts forming. A weak solution of baby shampoo in warm water is ideal.
- **A small operation** is an option if it is troublesome or persistent. Your GP can refer you to an eye surgeon (ophthalmologist) for this. The operation is usually done under local anaesthetic, although children and some adults may not tolerate this and may require general anaesthetic. The eyelid is numbed. A small cut is then made on the inside of the eyelid to release the contents of the cyst and it is scraped out. After surgery antibiotic drops or ointment are commonly prescribed.

Are there any complications?

Most chalazia cause no problems. Rarely, a cyst can become infected and this infection can spread to involve the whole eyelid and tissues surrounding the eye. The eyelid may be very swollen and red. You might not be able to open the eye and you may have a lot of pain and a high temperature (fever). If you develop this type of complication, called orbital cellulitis, you need to see a medical professional urgently. Treatment of orbital cellulitis is with antibiotics, usually into a vein (intravenously) via a drip, in hospital.

Will it happen again?

For most people a chalazion occurs just once. However, some people are prone to developing them and they may come back (recur). You may be able to prevent this by using a hot compress on the eyelids (described above) and massaging the eyelids each morning.

Further reading & references

- [Arbabi EM, Kelly RJ, Carrim ZI; Chalazion. BMJ. 2010 Aug 10;341:c4044. doi: 10.1136/bmj.c4044.](#)
- [Lindsley K, Nichols JJ, Dickersin K; Interventions for acute internal hordeolum. Cochrane Database Syst Rev. 2013 Apr 30;4:CD007742. doi: 10.1002/14651858.CD007742.pub3.](#)
- [Meibomian cyst \(chalazion\); NICE CKS, November 2015 \(UK access only\)](#)

Disclaimer: This article is for information only and should not be used for the diagnosis or treatment of medical conditions. Patient Platform Limited has used all reasonable care in compiling the information but makes no warranty as to its accuracy. Consult a doctor or other healthcare professional for diagnosis and treatment of medical conditions. For details see our [conditions](#).

| | | |
|----------------------------|---------------------------------|----------------------------|
| Author: Dr Mary Lowth | Peer Reviewer: Dr Colin Tidy | |
| Document ID: 4570 (v40) | Last Checked: 10/11/2016 | Next Review: 10/11/2019 |

View this article online at: patient.info/chalazion

Discuss Chalazion and find more trusted resources at [Patient](#).

Ask your doctor about Patient Access

- 🔍 Book appointments
- 🔍 Order repeat prescriptions
- 🔍 View your medical record
- 🔍 Create a personal health record (iOS only)



Simple, quick and convenient.
Visit patient.info/patient-access
or search 'Patient Access'