Health and care working together in South Yorkshire and Bassetlaw

Tonsillectomy – Prior Approval Through IFR

Special ity	Procedure	Criteria for treatment	Evidence Base	Process	Date of review
ENT	Tonsillectomy	The CCG will only fund tonsillectomy when one or more of the following criteria have been met: Recurrent attacks of tonsillitis as defined by: Sore throats are due to acute tonsillitis which is disabling and prevents normal functioning AND 7 or more well documented, clinically significant *, adequately treated episodes in the preceding year OR 5 or more such episodes in each of the preceding 2 years OR 3 or more such episodes in each of the preceding 3 years Two or more episodes of Quinsy (peritonsillar abscess) Severe halitosis secondary to tonsillar crypt debris Failure to thrive secondary to difficulty swallowing caused by enlarged tonsils Sleep disordered breathing or obstructive sleep apnoea diagnosed by an overnight pulse oximetry or polysonography Biopsy/removal of lesion on tonsil A Clinically significant episode is characterised by at least one of the following: Oral temperature of at least 38.30C requiring antibiotic treatment Tender anterior cervical lymph nodes. Tonsillar exudates.	Burton MJ, Glasziou PP. Tonsillectomy or adenotonsillectomy versus non-surgical treatment for chronic/recurrent acute tonsillitis. Cochrane Database of Systematic Reviews 1999, Issue 3. Art. No.: CD001802. First published online: July 26 1999. Available from: http://www.cochrane.org/reviews/en/ab001802.html (accessed 2016) Paradise JL, Bluestone CD, Bachman RZ. Efficacy of tonsillectomy for recurrent throat infection in severely affected children. Results of parallel randomized and non-randomized clinical trials. N England J Med 1984:310(11):674-83 SIGN. Management of sore throat and indications for tonsillectomy. A National clinical Guideline. April 2010 http://www.sign.ac.uk/pdf/sign117.pd f (accessed 2016)	Barnsley CCG require prior approval through IFR for this procedure	Dec 2018