

View this article online at: patient.info/vaginal-dryness-atrophic-vaginitis

Vaginal Dryness (Atrophic Vaginitis)

Many women notice changes in their vagina and genital area after the menopause. These changes may include dryness and discomfort during sex.

There may also be bladder symptoms. These can all usually be improved with treatment. Treatment options include lubricating gels, hormone replacement therapy (HRT) or vaginal oestrogen preparations.

What is vaginal dryness and what causes it?

Before the menopause (often called the change of life) the skin and tissues around your vagina are kept supple and moist by fluids and mucus. These are made by glands at the neck of your womb. The female hormone, oestrogen, affects these glands. Oestrogen also affects your tissues in and around your vagina, causing the lining of your vagina to be thicker and more elastic. Oestrogen stimulates the cells that line your vagina to produce glycogen. Glycogen is a compound which encourages the presence of helpful germs (bacteria) which protect your vagina from infections.

After the menopause your ovaries make less oestrogen. The lack of oestrogen leads to thinning of the tissues around your vagina and a reduction in the number of glands that make mucus. You may also lose some fat tissue from around the genital area. This may make the area also look slightly different to how it was before the menopause.

In summary, the hormonal changes that occur during the menopause make your vagina shorter, less elastic and drier and may also affect your bladder. These changes usually take months or years to develop after the menopause and vary from woman to woman. Atrophic vaginitis is the medical term for the condition when these changes produce troublesome symptoms.

How common is vaginal dryness?

After the menopause at least half of women have some symptoms related to vaginal dryness. You are also more likely to experience symptoms as more years pass after your menopause. It is probably even more common than that, as many women are embarrassed or feel they do not want to trouble their doctor with these symptoms.

What symptoms can occur?

The changes described above can occur without causing any symptoms or discomfort. However, some of the following symptoms may develop in some women. Vaginal dryness is a common (and usually treatable) cause of the following problems. However, these problems can also be caused by other medical conditions.

- Pain when you have sex. This may occur because your vagina is smaller, drier and less likely to become lubricated during
 sex compared with how it was before the menopause. Also, the skin around your vagina is more fragile and this can make the
 problem worse.
- Discomfort if your vulva or vagina is sore and red.
- Vaginal discharge. There may be a white or yellow discharge. Sometimes this is due to an infection. Infection is more likely if the discharge is smelly and unpleasant.
- Itch. The skin around your vagina is more sensitive and more likely to itch. This can make you prone to scratching, which
 then makes your skin more likely to itch, and so on. This is called an itch/scratch cycle which can become difficult to break
 and can be distressing.
- Urinary problems. Vaginal dryness may contribute to various urinary problems. This is because of thinning and weakening of
 the tissues around the neck of your bladder, or around the opening for urine to pass (the urethra). For example, urinary
 symptoms that may occur include an urgency to get to the toilet and recurring urinary infections.

What are the treatments for vaginal dryness?

Not all women have all of the above symptoms. Treatment usually depends on which symptoms are the most troublesome. Because the problem is mainly due to a lack of oestrogen, it can often be helped by replacing the oestrogen in your tissues.

Oestrogen creams and other topical preparations

A cream, vaginal tablet or ring containing oestrogen is often prescribed. A vaginal tablet is a very small tablet that you insert into your vagina with a small applicator. The ring is a soft, flexible ring, 55 mm across, with a centre that contains the oestrogen hormone. This ring releases a steady, low dose of oestrogen each day and it lasts for three months.

These preparations work to restore oestrogen to your vagina and surrounding tissues without giving oestrogen to the whole body. Usually the treatment is used every day for about two weeks, and then twice a week for as long as is needed. This treatment usually works well but the symptoms may come back sometime after stopping the treatment. Occasionally a repeated course of using it daily for two weeks is needed. These preparations should not be used as additional lubrication during sex; lubricating gels should be used instead.

Note: the oestrogen creams may damage latex condoms and diaphragms; if you are using these types of contraception then it would be preferable either to use vaginal tablets or the vaginal ring.

Hormone replacement therapy (HRT)

This means taking oestrogen in the form of a tablet, gel or patches. This is often the best treatment for relieving your symptoms, especially if you are experiencing other symptoms of the menopause. There are advantages and disadvantages of using HRT. Read more about hormone replacement therapy.

Vaginal lubricants and moisturisers

If vaginal dryness is the only problem, or hormone creams are not recommended because of other medical problems, lubricating gels or moisturisers may help. There are different lubricants which can improve the dryness during sexual intercourse. You can buy these from the pharmacy and your pharmacist should be able to advise you.

Vaginal moisturisers can work really well to improve the moisture in your vagina. These need to be used regularly.

Note: Vaseline® is NOT recommended as a lubricant. It is not smooth or slippery enough and it can break down the latex in condoms.

Your symptoms should improve after about three weeks of treatment. You should see your doctor if your symptoms do not improve, as sometimes these symptoms can be due to other conditions. It is also very important to see your doctor if you have any bleeding from your vagina if you are receiving hormone treatment.

Further reading & references

- Panay N et al; British Menopause Society & Women's Heath Concern recommendations on hormone replacement therapy, May 2013
- Menopause: diagnosis and management; NICE Guidelines (Nov 2015)
- Kaunitz AM, Manson JE; Management of Menopausal Symptoms. Obstet Gynecol. 2015 Oct; 126(4):859-76. doi: 10.1097/AOG.0000000000001058.
- No authors listed; Nonhormonal management of menopause-associated vasomotor symptoms: 2015 position statement of The North American Menopause Society. Menopause. 2015 Nov;22(11):1155-74. doi: 10.1097/GME.00000000000546.
- Contraception for Women Aged over 40 Years; Faculty of Sexual and Reproductive Healthcare (August 2017)
- Langer RD; The evidence base for HRT: what can we believe? Climacteric. 2017 Apr;20(2):91-96. doi: 10.1080/13697137.2017.1280251. Epub 2017 Mar 10.
- Information for women with latrogenic Premature Ovarian Insufficiency, European Society of Human Reproduction and Embryology, 2016

Disclaimer: This article is for information only and should not be used for the diagnosis or treatment of medical conditions. Patient Platform Limited has used all reasonable care in compiling the information but makes no warranty as to its accuracy. Consult a doctor or other healthcare professional for diagnosis and treatment of medical conditions. For details see our conditions.

Author: Dr Jacqueline Payne	Peer Reviewer: Dr Sarah Jarvis MBE	
Document ID:	Last Checked:	Next Review:
4199 (v42)	26/02/2016	25/02/2019

View this article online at: patient.info/vaginal-dryness-atrophic-vaginitis

Discuss Vaginal Dryness (Atrophic Vaginitis) and find more trusted resources at Patient.





