

Sexually Transmitted Infections & HIV in Primary Care

Dr S Bates

Gateway Clinic, Sackville street

Our Service

- July 2015 – HIV and Sexual health care delivered from Gateway Clinic
- HIV care provided by Barnsley Hospital
- Contraception and STI care provided by Spectrum CIC

GUM Services

- Confidential
- Non judgemental
- Open access
- Onsite diagnosis
- Free treatment



"Ah, Mr Bond, I've been expecting you..."

Services delivered

Contraceptive pills

Contraceptive patches

Coils

Implants

**Assessment and referral for
abortion**

Psychosexual counselling

Youth clinic

Non-contraceptive use of IUS

Non-invasive screening

(asymptomatic)

Chlamydia screening

Point of care tests for HIV

Diagnosis and treatment of STI

Biopsy to confirm diagnosis

Motivational interviewing

Contact tracing

**Hepatitis B vaccination (sexual
risk)**

Genital dematoses

**Non-STI causes of vaginal
discharge**

How to refer

- Allow patients to self refer and contact the clinic directly
 - Patient may not make contact
- 'Phone the clinic to make an appointment.
 - Ask for health advisors if confirmed STI or complex
- Refer by letter
 - delay of receiving an appointment.

Contact details

Gateway Clinic

Gateway Plaza

Sackville Street

Barnsley

S70 2RD

Tel: 0800 055 6442

STIs- general principles

- Accurate diagnosis
- Prompt, appropriate and simple treatment regimens
- Contact tracing
- Test of cure where appropriate

Infection screening


When to think about Screening

- PCB / IMB
- Cervicitis
- Abnormal vaginal discharge
- Genital ulceration
- Dysuria
- Rectal / anal pain / discharge
- Skin rashes
- Evidence on a co-existing STI
- Recent partner change or high risk group / partner

Tests – CT GC NAAT

- Men
 - 1st void urine (must have held urine for 1 hour)
- Women
 - Endo-cervical swab
 - Self or physician taken vulvo vaginal swabs

Gonorrhoea - testing



Caution

- A positive test does NOT mean an individual has gonorrhoea
- Patients with a positive NAAT test must have culture tests to confirm diagnosis
- Preferably should be referred to GUM



Chlamydia and gonorrhoea – Treatment

Chlamydia


- Azithromycin 1 gr PO single dose
- Doxycycline 100mg bd 1/52 for rectal chlamydia

Consider referral to GUM for contact tracing

Gonorrhoea

Refer to GUM

- Ceftriaxone 500mg IM single dose with azithromycin 1 gram PO single dose



Genital blisters / ulcers

HSV

Treatment:
Valaciclovir
500mg BD
5/7

Refer to

GUM
Tests!

HSV PCR from
lesion




Syphilis

If suspicious of syphilis
refer to GUM

Treatment: Benzathine
penicillin 2.4MU IM for
primary



STS - Serological
tests for syphilis
** may be
negative in early
disease



Dark ground
microscopy





Genital warts

HPV Types 6 & 11 account for 90% of genital warts

HPV 6 & 11 infections are usually asymptomatic and resolve spontaneously - 90% within two years.



Treatment

- Destructive methods
 - LN2
 - podophylotoxin
 - Surgery
 - Diathermy
 - Curettage
- Immune response modifiers
 - Imiquimod

What's new?

- Resistant GC
- Chem sex
- LGV
- Hepatitis C

LGV

- Chlamydia – invasive serovars L1,L2,L3
 - 80 cases per quarter
 - UK – highest global rates in MSM
 - 77% London, Brighton, Manchester
- Symptoms
 - Primary
 - Painless papule / pustule – may persist for weeks
 - Proctitis (96%) – blood/pus/pain
 - Secondary
 - Lymphadenopathy / bubo formation, usually unilateral ‘groove’ sign
 - Fever, arthritis, pneumonitis, hepatitis




- Diagnosis
 - NAAT testing for CT, if positive test for LGV serovars at Colindale
- Treatment
 - 3 weeks of doxycycline 100 mg BD (or erythromycin / azithromycin)

HIV


Disease process,
Epidemiology & Treatment

1. How many people are estimated to be living with HIV in the UK?

- A. <50,000
- B. 50-75,000
- C. 75,-100,000
- D. >100,000




How many people are estimated to be living with HIV in the UK?


- A. <50,000
- B. 50 - 75,000
- C. 75- 100,000 
- D. >100,000

2. What proportion of people with HIV remain undiagnosed?

- A. <10 %
- B. 10-20%
- C. 20-30%
- D. 30-40%
- E. >40%




What proportion of people with HIV remain undiagnosed?

- A. <10 %
- B. 10-20%
- C. 20-30%
- D. 30-40% 
- E. >40%

3. How many people living with HIV are accessing services in Barnsley?

- A. <50
- B. 50-100
- C. 100-150
- D. 150-200
- E. >200




How many people living with HIV are accessing services in Barnsley?

- A. <50
- B. 50-100
- C. 100-150
- D. 150-200
- E. >200

4. What is the main cause of death in people living with HIV?

- A. Conditions unrelated to HIV
- B. Treatment failure
- C. Late diagnosis of infection
- D. Pneumocystis jirovecii




What is the main cause of death in people living with HIV?

- A. Conditions unrelated to HIV
- B. Treatment failure ✓
- C. Late diagnosis of infection
- D. *Pneumocystis jirovecii*


5. In order to do an HIV tests, pre-test counselling has to be undertaken

- A. True
- B. False




6. It takes 3 months following exposure for HIV to be reliably diagnosed on a blood test

- A. True
- B. False




7. HIV tests should only be requested by practitioners with experience in sexual health issues

- A. True
- B. False



8. Any practitioner can request an HIV test, with verbal consent from the patient.

- A. True
- B. False



Which of the following statements is true?

- A. In order to do an HIV tests, pre-test counselling has to be undertaken
- B. It takes 3 months following exposure for HIV to be reliably diagnosed on a blood test
- C. HIV tests should only be requested by practitioners with experience in sexual health issues
- D. Any practitioner can request an HIV test, with verbal consent from the patient.

HIV

- the basics

Acquisition

- Sex
- Vertical transmission
- Injecting drug use
- Blood transfusions with infected blood
- Percutaneous injury – needle stick

HIV infection

Acute infection -
seroconversion



Asymptomatic



HIV related
illnesses



AIDS defining



illness

Sero-conversion	CD4 > 500	CD4 500 - 200	CD4 <200
<ul style="list-style-type: none"> ○Fever ○Myalgia ○Arthralgia ○Adenopathy ○Malaise ○Rash ○Meningo-encephalitis 	<ul style="list-style-type: none"> ○Guillain-Barre syndrome ○Bell's palsy ○Polymyositis ○Chronic demyelinating neuropathy ○Idiopathic thrombocytopenia ○Tinea 	<ul style="list-style-type: none"> ○Seborrhoeic dermatitis ○Gingivitis ○Warts ○Molluscum ○TB ○Herpes Zoster/Simplex ○Oral candida ○KS ○CIN ○Primary CNS lymphoma 	<ul style="list-style-type: none"> ○Cryptosporidiosis ○PCP ○Toxoplasmosis ○Cryptococcal meningitis ○CMV ○MAC

Primary HIV infection

Classical triad of:

- Pharyngitis

- Rash

- Fever

Clinical Manifestations





















HIV in the UK

- There are currently an estimated 96,000 people with HIV in the UK.
- 24% of those infected (>23,000) remain undiagnosed – HPA 2011
- 47% of new HIV diagnoses are diagnosed late (CD4 < 350). 56% of women are diagnosed late.
- 35% of HIV-related deaths attributable to late diagnosis – BHIVA audit 2006
- Effective treatments greatly reduce morbidity
- Better response to treatment if started early
- Fully suppressed viral loads reduce onward transmission

Estimated number of adults (15 to 59 years) living with HIV (both diagnosed and undiagnosed) in the UK: 2007



When to test?



TOP 3

1. TB
2. BBV – Hep B / C
3. Lymphoma

B. An HIV test should be considered in the following settings where diagnosed HIV prevalence in the local population (PCT/LA) exceeds 2 in 1000 population (see local PCT data[†]):

1. all men and women registering in general practice
2. all general medical admissions.

The introduction of universal HIV testing in these settings should be thoroughly evaluated for acceptability and feasibility and the resultant data made available to better inform the ongoing implementation of these guidelines.

C. HIV testing should be also routinely offered and recommended to the following patients:

1. all patients presenting for healthcare where HIV, including primary HIV infection, enters the differential diagnosis (see table of indicator diseases and section on primary HIV infection)
2. all patients diagnosed with a sexually transmitted infection
3. all sexual partners of men and women known to be HIV positive
4. all men who have disclosed sexual contact with other men
5. all female sexual contacts of men who have sex with men
6. all patients reporting a history of injecting drug use
7. all men and women known to be from a country of high HIV prevalence (>1%*)
8. all men and women who report sexual contact abroad or in the UK with individuals from countries of high HIV prevalence.*

* for an up to date list see

UK National Guidelines for HIV Testing 2008

Table 1: Clinical indicator diseases for adult HIV infection

	AIDS-defining conditions	Other conditions where HIV testing should be offered
Respiratory	Tuberculosis Pneumocystis	Bacterial pneumonia Aspergillosis
Neurology	Cerebral toxoplasmosis Primary cerebral lymphoma Cryptococcal meningitis Progressive multifocal leucoencephalopathy	Aseptic meningitis/encephalitis Cerebral abscess Space occupying lesion of unknown cause Guillain–Barré syndrome Transverse myelitis Peripheral neuropathy Dementia Leucoencephalopathy
Dermatology	Kaposi's sarcoma	Severe or recalcitrant seborrhoeic dermatitis Severe or recalcitrant psoriasis Multidermatomal or recurrent herpes zoster

Gastroenterology	Persistent cryptosporidiosis	Oral candidiasis Oral hairy leukoplakia Chronic diarrhoea of unknown cause Weight loss of unknown cause Salmonella, shigella or campylobacter Hepatitis B infection Hepatitis C infection
Oncology	Non-Hodgkin's lymphoma	Anal cancer or anal intraepithelial dysplasia Lung cancer Seminoma Head and neck cancer Hodgkin's lymphoma Castleman's disease
Gynaecology	Cervical cancer	Vaginal intraepithelial neoplasia Cervical intraepithelial neoplasia Grade 2 or above
Haematology		Any unexplained blood dyscrasia including: <ul style="list-style-type: none">• thrombocytopenia• neutropenia• lymphopenia
Ophthalmology	Cytomegalovirus retinitis	Infective retinal diseases including herpesviruses and toxoplasma

Haematology

Any unexplained blood dyscrasia including:

- thrombocytopenia
- neutropenia
- lymphopenia

Ophthalmology

Cytomegalovirus retinitis

Infective retinal diseases including herpesviruses and toxoplasma

Any unexplained retinopathy

ENT

Lymphadenopathy of unknown cause

Chronic parotitis

Lymphoepithelial parotid cysts

Other

Mononucleosis-like syndrome (primary HIV infection)

Pyrexia of unknown origin

Any lymphadenopathy of unknown cause

Any sexually transmitted infection

HAART

- At least 4 useful, widely available single tablet regimens
- ‘Cleaner’ drugs – fewer side effects
- Newer drug groups with fewer drug-drug interactions
- Compliance is the key to successful treatment
ART is for life - *Never* advise a patient to stop ART without advice from HIV unit



Benefits of treatment

- Preservation of immune function
- Improved life expectancy
- Reduced onward transmission
- Reduction of mother to baby transmission
- Fewer hospital admissions

What to watch out for in Primary Care

- A few patients still choose not to disclose
- Watch out for drug interactions. Common culprits:
 - Statins
 - Omeprazole

www.hiv-druginteractions.org

- Live vaccinations

How to do an HIV test

You need one of these....



.....verbal consent and a standard microbiology form – simply request HIV test!

You do not need....

Formal pre test counselling*, written consent, cat 3 stickers (unless other risk), special permission!

** If patient identified as being at high risk of infection GUM health advisors will come and see patient*



